

LICHENS AND AIR QUALITY
IN
WHITE MOUNTAIN NATIONAL FOREST
WILDERNESS AREAS

Final Report

U. S. Forest Service
Contract USDA 40-1484-7-614

by

Clifford M. Wetmore
Botany Department
University of Minnesota
St. Paul, Minnesota

June, 1989

TABLE OF CONTENTS

LICHENS OF WHITE MOUNTAINS WILDERNESS AREAS

	Page
Preface.....	1
Introduction.....	2
Methods.....	5
Lichen Flora.....	6
Species List.....	7
Discussion of the Lichen Flora.....	11
Elemental analysis.....	14
Methods.....	14
Results and Discussion.....	16
Conclusions.....	17
Literature Cited.....	18
Appendix I: Collection Localities.....	27
Map of Collection Localities	
Appendix II: Species Sensitive to Sulphur Dioxide.....	30
Maps of Sensitive Species	
Appendix III: Species Cited in Literature from White Mts...	31

PREFACE

Under a grant from the U. S. Forest Service a lichen study was to be performed in the Presidential Dry River and the Great Gulf Wilderness Areas of the White Mountain National Forest. This study was to survey the lichens of the areas, produce a lichen flora, collect and analyze lichens for chemical contents and evaluate the lichen flora with reference to the air quality. This study is to establish baseline data for future restudy and determine the presence of any air quality problems as might be shown by the lichens at the time of the study. All work was done at the University of Minnesota with consultation with Randy Ferrin, Laconia, NH and with other U. S. Forest Service personnel.

The forest personnel have been very helpful during the field work with assistance and transportation which has contributed significantly to the success of the project. The study was made possible by funds from the U. S. Forest Service. The assistance of all of these is gratefully acknowledged.

INTRODUCTION

Lichens are composite plants composed of two different types of organisms. The lichen plant body (thallus) is made of fungi and algae living together in a symbiotic arrangement in which both partners are benefited and the composite plant body can grow in places where neither component could live alone. The thallus has no protective layer on the outside, such as the epidermis of a leaf, so the air in the thallus has free exchange with the atmosphere. Lichens are slow growing (a few millimeters per year) and remain alive for many years and so must have a habitat that is relatively undisturbed in order to survive. Lichens vary greatly in their ecological requirements but almost all of them can grow in places that only receive periodic moisture. When moisture is lacking they go dormant until the next rain or dew-fall. Some species can grow in habitats with very infrequent occurrences of moisture while others need high humidity and frequent wetting in order to survive. This difference in moisture requirements is very important in the distribution of lichens.

Lichens are known to be very sensitive to low levels of many atmospheric pollutants. Many are damaged or killed by levels of sulfur dioxide, nitrogen oxides, fluorides or ozone alone or in various combinations. Levels of sulfur dioxide as low as 13 ug/cubic meter (annual average) will cause the death of some lichens (LeBlanc et al., 1972). Other lichens are less sensitive and a few can tolerate levels of sulfur dioxide over 300 ug/cubic meter (Laundon, 1967, Trass, 1973). The algae of the thallus are the first to be damaged in areas with air pollution and the first indication of damage is discoloring and death of the algae causing bleached lobes, which quickly leads to the death of the lichen. After the lichen dies it disappears from the substrate within a few months to a year as it disintegrates and decomposes (Wetmore, 1982).

Lichens are more sensitive to air pollution when they are wet and physiologically active and are least sensitive when dry (Nash, 1973, Marsh & Nash, 1979) and are more sensitive when growing on acid substrates.

Contrary to some published reports (Medlin, 1985) there is little evidence that most lichens are good indicators of acid precipitation. However, Sigal & Johnston (1986) have reported that one

species of *Umbilicaria* shows visible damage due to artificial acid rain. They also report that similar symptoms were found in collections from various localities in North America. Lechowicz (1987) reported that acid rain only slightly reduced growth of *Cladina stellaris* but Hutchinson et al. (1986) reported that extremely acid precipitation killed or damaged some mosses and lichens. Scott & Hutchinson (1987) showed temporary reduction of photosynthesis in *Cladina stellaris* and *C. rangiferina* after artificial acid rain.

Lichens are able to accumulate chemical elements in excess of their metabolic needs depending on the levels in the substrate and the air and, since lichens are slow growing and long lived, they serve as good summarizers of the environmental conditions in which they are growing. Chemical analysis of the thallus of lichens growing in areas of high fallout of certain elements will show elevated levels in the thallus. Toxic substances (such as sulfur) are also accumulated and determination of the levels of these toxic elements can provide indications of the sub-lethal but elevated levels in the air.

The Great Gulf Wilderness Area is located just north of Mt. Washington in northern New Hampshire and the Presidential Dry River Wilderness Area is located just south of Mt. Washington. Only these two wilderness areas in the White Mts. National Forest were included in this study and Mt. Washington itself was outside of both wilderness areas. The elevations range from 1200 ft to over 5300 ft. The terrain is steep and rugged with granitic peaks and ridges. The forests at the lower elevations are mixed hardwoods with sugar maple (*Acer saccharum*), beech (*Fagus grandifolia*), yellow birch (*Betula allegheniensis*), white birch (*Betula papyrifera*) and some areas of hemlock (*Tsuga canadensis*) and white pine (*Pinus strobus*). At higher elevations there are balsam fir (*Abies balsamifera*) and black spruce (*Picea mariana*) grading into krumholz at the highest elevations. On the highest peaks alpine vegetation occurs among the abundant rocks.

The White Mountains were one of the first areas of North America to be extensively studied by Edward Tuckerman and his co-workers in their lichenological studies over 150 years ago. These early lichenologists thoroughly collected around Mt. Washington and throughout the White Mts. for decades during the 19th century. Tuckerman published many reports on the lichens of the region

(see literature cited). Until recent times this region was probably one of the best known areas of North America lichenologically. Since Tuckerman's time most North American lichenologists have collected around Mt. Washington at some time or another. Many ecologists have studied the alpine vegetation around Mt. Washington and some studies included lichens in their papers (Bliss, 1962, 1963, 1966, Bliss & Hadley, 1964, Hadley & Bliss, 1964). Many of the lichen collections of various collectors have been cited in monographs and revisions and these references are listed in the literature cited. A list of the 289 lichens reported in the literature is included in Appendix III. The names in Appendix III have been brought up to current nomenclatural practice according to Egan (1987) wherever possible but some of the smaller genera used by Egan are not recognized here.

METHODS

Field work was done during July, 1988 when 1005 collections were made at 21 localities. A complete list of collection localities is given in Appendix I and are indicated on Fig. 1. Localities for collecting were selected first to give a general coverage of the wilderness areas and be within practical walking distance from roads, second, to sample all vegetational types, and third, to be in localities that should be rich in lichens. Undisturbed as well as disturbed habitats were studied. At each locality voucher specimens of all species found were collected to record the total flora for each locality and to avoid missing different species that might appear similar in the field. At some localities additional material of selected species was collected for chemical analysis (see below). While collecting at each locality observations were made about the general health of the lichens.

Identifications were carried out at the University of Minnesota with the aid of comparison material in the herbarium and using thin layer chromatography for identification of the lichen substances where necessary. The original packet of each collection has been deposited in the University of Minnesota Herbarium. All specimens deposited at the University of Minnesota have been entered into the herbarium computerized data base maintained there. Lists of species found at each locality are available from this data base at any time on request.

LICHEN FLORA

The following list of lichens is based only on my collections. The species reported in the

literature are given in Appendix III. Species found only once are indicated by "Rare". In the first columns the letters indicate the sensitivity to sulfur dioxide, if known, according to the categories proposed by Wetmore (1983): S=Sensitive, I=Intermediate, T=Tolerant. SI is intermediate between Sensitive and Intermediate and I-T is intermediate between Intermediate and Tolerant. Species in the Sensitive category are absent when annual average levels of sulfur dioxide are above 50ug per cubic meter. The Intermediate category includes those species present between 50 and 100ug and those in the Tolerant category are present at over 100ug per cubic meter.

SPECIES LIST

- Acarospora fuscata (Nyl.) Arn.
- Acarospora sinopica (Wahlenb. in Ach.) K"orb.
- Arthothelium ruanum (Mass.) Zw.
1 additional unidentified species of Arthothelium
- Aspicilia caesiocinerea (Nyl. ex Malbr.) Arn. Rare
- Aspicilia cinerea (L.) K"orb.
- Bacidia chlorantha (Tuck.) Fink
- Bacidia lugubris (Somm.) Zahlbr. Rare
- Bacidia schweinitzii (Tuck.) Schneid. Rare
1 additional unidentified species of Bacidia
- Baeomyces rufus (Huds.) Rebent.
- Botrydina viridis (Ach.) Redh. & Kuyper
- SBryoria furcellata (Fr.) Brodo & Hawksw.
- IBryoria fuscescens (Gyeln.) Brodo & Hawksw.
- Bryoria nadvornikiana (Gyeln.) Brodo & Hawksw.
- Buellia arnoldii Serv.
- Buellia disciformis (Fr.) Mudd
- IBuellia stillingiana Steiner
- Calicium trabinellum (Ach.) Ach.
- S-ICaloplaca cerina (Ehrh. ex Hedw.) Th. Fr. Rare
- ICaloplaca holocarpa (Hoffm.) Wade
- S-ICetraria ciliaris Ach.
- Cetraria cucullata (Bellardi) Ach.
- Cetraria delisei (Bory ex Schaer.) Nyl.
- Cetraria halei W. & C. Culb.
- Cetraria hepatizon (Ach.) Vain.
- Cetraria islandica (L.) Ach.
- Cetraria nivalis (L.) Ach.
- Cetraria oakesiana Tuck.
- ICetraria orbata (Nyl.) Fink
- ICetraria pinastri (Scop.) Gray
- ICetraria sepincola (Ehrh.) Ach.
- Cetrelia cetrariooides (Del. ex Duby) W. & C. Culb. Rare
- Cetrelia olivetorum (Nyl.) W. & C. Culb.
- Chaenotheca brunneola (Ach.) M"ull. Arg.
- Chaenotheca xylooxena Nadv. Rare
- Chaenothecopsis debilis (Turn. & Borr. ex Sm.) Tibell
- Cladina arbuscula (Wallr.) Hale & W. Culb.

Cladina mitis (Sandst.) Hustich
Cladina rangiferina (L.) Nyl.
Cladina stellaris (Opiz) Brodo
Cladina stygia (Fr.) Ahti
Cladonia amaurocraea (Fl"orke) Schaer.
Cladonia bacillaris Nyl.
Cladonia caespiticia (Pers.) Fl"orke
Cladonia cenotea (Ach.) Schaer. Rare
Cladonia chlorophaea (Fl"orke ex Somm.) Spreng.
Cladonia coccifera (L.) Willd.
ICladonia coniocraea (Fl"orke) Spreng.
Cladonia cornuta (L.) Hoffm.
Cladonia crispata (Ach.) Flot.
ICladonia cristatella Tuck.
Cladonia cyanipes (Somm.) Nyl.
Cladonia deformis (L.) Hoffm.
Cladonia digitata (L.) Hoffm.
S-ICladonia fimbriata (L.) Fr. Rare
Cladonia floerkeana (Fr.) Fl"orke
Cladonia furcata (Huds.) Schrad.
Cladonia gracilis (L.) Willd.
Cladonia macilenta Hoffm. Rare
Cladonia maxima (Asah.) Ahti Rare
Cladonia merochlorophaea Asah.
Cladonia multiformis G. K. Merr. Rare
Cladonia phyllophora Ehrh. ex Hoffm. Rare
Cladonia pleurota (Fl"orke) Schaer.
Cladonia pseudorangiformis Asah.
Cladonia pyxidata (L.) Hoffm. Rare
Cladonia scabriuscula (Del.in Duby) Leight. Rare
Cladonia squamosa (Scop.) Hoffm.
Cladonia subsulculata Nyl. Rare
Cladonia sulphurina (Michx.) Fr.
Cladonia uncialis (L.) Web. ex Wigg.
Cladonia verticillata (Hoffm.) Schaer. Rare
Conotrema urceolatum (Ach.) Tuck.
Cornicularia aculeata (Schreb.) Ach.
Dimelaena oreina (Ach.) Norm. Rare
Diploschistes scruposus (Schreb.) Norm.
IEvernia mesomorpha Nyl.
IGraphis scripta (L.) Ach.
Haematomma cismonicum Beltr.
Haematomma elatinum (Ach.) Mass.
Haematomma lapponicum R"as.
IHaematomma ochrophaeum (Tuck.) Mass.
Hymenelia lacustris (With.) Poelt & Vezda Rare
Hypogymnia krogiae Ohls.
IHypogymnia physodes (L.) Nyl.
IHypogymnia tubulosa (Schaer.) Hav.
Icmadophila ericetorum (L.) Zahlbr.
Ilmshaugia aleurites (Ach.) S. F. Meyer
Julella fallaciosa (Stizenb. ex Arn.) R. Harris
Lasallia papulosa (Ach.) Llano
Lecanora badia (Hoffm.) Ach.

Lecanora caesiorubella Ach. subsp. caesiorubella
IILecanora circumborealis Brodo & Vitik.
Lecanora glabrata (Ach.) Malme
Lecanora hypocarpa (Tuck.) Brodo
Lecanora mutabilis Somm.
Lecanora piniperda K"orb. Rare
Lecanora polytropa (Hoffm.) Rabenh.
IILecanora pulicaris (Pers.) Ach.
IILecanora symmicta (Ach.) Ach. Rare
Lecanora thysanophora Harris ined.
Lecanora wisconsinensis Magn.
ILecidea albofuscens Nyl. Rare
Lecidea albohyalina (Nyl.) Th. Fr.
Lecidea diapensiae Th. Fr.
Lecidea epixanthoidiza Nyl.
Lecidea helvola (K"orb. ex Hellb.) Oliv.
Lecidea plana (Lahm in K"orb.) Nyl.
Lecidea tessellata Fl"orke Rare
Lecidea tornoensis Nyl.
S Lecidea vernalis (L.) Ach.
2 additional unidentified species of Lecidea
Lecidella euphorea (Fl"orke) Hert.
Lecidella stigmatea (Ach.) Hert. & Leuck.
Lecidoma demissum (Rutstr.) Schneid. & Hert.
Lepraria finkii (B. de Lesd. in Hue) R. Harris
Lepraria neglecta (Nyl.) Lett.
Leptogium cyanescens (Rabenh.) K"orb.
Leptorraphis epidermidis (Ach.) Th. Fr.
SLobaria pulmonaria (L.) Hoffm.
Lobaria quercizans Michx.
IILopadium pezizoideum (Ach.) K"orb.
Micarea bauschiana (K"orb.) V. Wirth & Vezda Rare
Micarea lignaria (Ach.) Hidl.
Micarea melaena (Nyl.) Hidl.
Micarea peliocarpa (Anzi) Coppins & R. Sant.
Micarea prasina Fr.
Micarea turfosa (Mass.) Du Rietz Rare
2 additional unidentified species of Micarea
IMycoblastus sanguinarius (L.) Norm.
Mycocalicium subtile (Pers.) Szat.
Nephroma parile (Ach.) Ach.
Nephroma resupinatum (L.) Ach. Rare
SOchrolechia androgyna (Hoffm.) Arn.
Ochrolechia arborea (Kreyer) Almb.
Ochrolechia frigida (Sw.) Lynge
Ochrolechia pallescens (L.) Mass.
Ochrolechia pseudopallescens Brodo Rare
Orphniospora moriopsis (Mass.) D. Hawksw.
Parmelia aurulenta Tuck.
Parmelia baltimorensis Gyeln.
Parmelia centrifuga (L.) Ach.
Parmelia cumberlandia (Gyeln.) Hale Rare
Parmelia galbina Ach. Rare
Parmelia halei Ahti

Parmelia omphalodes (L.) Ach.
Parmelia panniformis (Nyl.) Vain.
IParmelia rupestris Ach.
IParmelia saxatilis (L.) Ach.
IParmelia septentrionalis (Lynge) Ahti Rare
SParmelia squarrosa Hale
Parmelia stygia (L.) Ach.
S-I Parmelia subaurifera Nyl.
I-TParmelia sulcata Tayl.
1 additional unidentified species of Parmelia
IParmeliopsis ambigua (Wulf. in Jacq.) Nyl.
IParmeliopsis hyperocea (Ach.) Arn.
Peltigera leucophlebia (Nyl.) Gyeln. Rare
Peltigera polydactyla (Neck.) Hoffm.
Peltigera praetextata (Fl"orke ex Somm.) Zopf Rare
IPertusaria amara (Ach.) Nyl.
Pertusaria consocians Dibb.
Pertusaria leucostoma (Bernh.) Mass. Rare
Pertusaria macounii (Lamb) Dibb. Rare
Pertusaria ophthalmiza (Nyl.) Nyl.
Pertusaria propinqua M"ull. Arg.
Pertusaria trachythallina Erichs. Rare
Pertusaria velata (Turn.) Nyl.
Pertusaria waghornei Hult. Rare
2 additional unidentified species of Pertusaria
IPhaeophyscia orbicularis (Neck.) Moberg Rare
Phaeophyscia rubropulchra (Degel.) Moberg
IPhyscia aipolia (Ehrh. ex Humb.) F"urnr. Rare
Placynthiella icmalea (Ach.) Coppins & James
Placynthiella oligotropha (Laund.) Coppins & James
Placynthiella uliginosa (Schrad.) Coppins & James
IPlatismatia glauca (L.) W. & C. Culb.
Platismatia tuckermanii (Oakes) W. & C. Culb. Rare
Porpidia crustulata (Ach.) Hert. & Knoph
Porpidia macrocarpa (DC. in Lam. & DC.) Hert. & Schwab
2 additional unidentified species of Porpidia
Pseudevernia cladonia (Tuck.) Hale & W. Culb.
Pseudevernia consocians (Vain.) Hale & W. Culb.
Pycnothelia papillaria Duf.
Pyrenula pseudobufonia (Rehm.) R. Harris
Pyxine sorediata (Ach.) Mont.
Ramalina intermedia (Del. ex Nyl.) Nyl. Rare
Rhizocarpon badioatrum (Fl"orke ex Spreng.) Th. Fr. Rare
Rhizocarpon cinereovirens (M"ull. Arg.) Vain.
Rhizocarpon eupetraeoides (Nyl.) Blomb. & Forss.
Rhizocarpon geographicum (L.) DC.
Rhizocarpon hochstetteri (K"orb.) Vain.
Rhizocarpon lecanorinum (K"orb.) Anders
Rhizocarpon obscuratum (Ach.) Mass. Rare
Rhizocarpon oederi (Weber) K"orb.
Rhizoplaca chrysoleuca (Sm.) Zopf Rare
IScoliosporum chlorococcum (Graewe ex Sten.) Vezda
Sphaerophorus fragilis (L.) Pers.
IStenocybe major Nyl. ex K"orb.

Stereocaulon dactylophyllum Fl"orke
Stereocaulon glaucescens Tuck.
Stereocaulon nanodes Tuck. Rare
Stereocaulon saxatile Magn.
Stereocaulon tomentosum Fr. Rare
1 additional unidentified species of Stereocaulon
Thamnolia subuliformis (Ehrh.) W. Culb.
Trapelia coarctata (Sm.) Choisy in Werner Rare
Trapelia involuta (Tayl.) Hert.
Trapeliopsis flexuosa (Fr.) Coppins & James Rare
Trapeliopsis gelatinosa Fl"orke) Coppins & James
Trapeliopsis granulosa (Hoffm.) Lumbsch.
Trapeliopsis viridescens (Schrad.) Coppins & James Rare
Umbilicaria deusta (L.) Baumg.
Umbilicaria hyperborea (Ach.) Hoffm.
Umbilicaria mammulata (Ach.) Tuck. Rare
Umbilicaria muehlenbergii (Ach.) Tuck.
Umbilicaria proboscidea (L.) Schrad.
Umbilicaria torrefacta (Lightf.) Schrad.
Umbilicaria vellea (L.) Ach.
SUsnea filipendula Stirt. Rare
S-IUsnea hirta (L.) Weber ex Wigg.

DISCUSSION OF FLORA

This list includes 214 species collected for this study. There are also an additional 12 unidentified species, some of which are undescribed. The most common species are Cetraria islandica, Cladina rangiferina, Cladina stellaris, Cladonia amaurocraea, Hypogymnia physodes, Mycoblastus sanguinarius, and Ochrolechia frigida.

Many alpine species occur at the higher elevations of the White Mts., such as Thamnolia vermicularis, Cetraria nivalis, and others. At lower elevations in the conifer forests the lichen flora is typical of the northern boreal forest that occurs across North America and includes Hypogymnia physodes, Cladina stellaris, Calicium spp. and others. A few species that are restricted to eastern North America also occur in the White Mts., such as Cladonia maxima, Hypogymnia krogiae, and Pseudevernia cladonia. In the hardwood forests there are fewer species than in the conifer forests, probably due to the dense shade.

Some of the rare species are rare wherever they are found throughout their distributional range and might be found at other localities with further searching and others may require special substrates that are rare in the area.

Some localities that should have many species do not. An example is near the Halfway House locality on the Mt. Washington Road. This balsam fir forest should have had many more species of lichens. It is probably not due to the dead and dying conifers (possibly due to acid precipitation) because this opens up the forest and provides dead branches for good lichen growth. Whether this lack of species is natural or real is unknown. Periodic restudy of the same areas might provide some answers.

Most of the historical species were found during this study but there are a number of species that might be expected but were not found. Some of the more remote localities within the wilderness areas may have produced additional species. One cannot make valid comparisons between the collections for this study and the historical species list. The length of time the many lichenologists in the past collected in the area compared to the length of time available for this study is not comparable. Many of the old records give the locality as "White Mts." which covers much more than the present wilderness areas. Many species in the literature were found in Tuckerman's Ravine and on Mt. Washington, which are outside of the present study areas. For these reasons it is impossible to attribute all of the missing species to air pollution. However, the northeastern United States is known to have severe air pollution problems and many of the missing species may no longer occur in the area.

Another way of analyzing the lichen flora of an area is to study the distributions of the sensitive species within the National Forest to look for voids in the distributions that might be caused by air pollution. Showman (1975) has described and used this technique in assessing sulfur dioxide levels around a power plant in Ohio. Only the very common species have meaning with such a technique since the rare species may be absent due to other factors.

Many of the lichens in the area have known sensitivity to sulfur dioxide according to the list presented in Wetmore (1983). Species in the most sensitive category are usually absent when sulfur dioxide levels are above 50ug per cubic meter average annual concentrations. The species that occur in the study area in the most sensitive category are as follows.

Bryoria furcellata

Lecidea vernalis

Lobaria pulmonaria

Ochrolechia androgyna

Parmelia squarrosa

Usnea filipendula

The distributions of these species are mapped Fig. 27. Although these species are not found at all localities and many are not common, there is no indication that the voids in the distributions are due to poor air quality. Some of the localities where collections were made do not have suitable habitats or substrates for some of these species. This is especially true for Lobaria pulmonaria that requires moist habitats.

There were no cases where lichens sensitive to sulfur dioxide were observed to be damaged or killed. All species normally found fertile were also fertile in the wilderness areas. These observations indicate that there probably is no local air quality degradation due to sulfur dioxide that causes observable damage to the lichen flora in the wilderness areas but the regional air quality may have had an effect on the lichen flora.

ELEMENTAL ANALYSIS

An important method of assessing the effects of air quality is by examining the elemental content of the lichens (Nieboer et al, 1972, 1977, 1978; Erdman & Gough, 1977; Puckett & Finegan, 1980; Nash & Sommerfeld, 1981). Elevated but sublethal levels of sulfur or other elements might indicate incipient damaging conditions.

METHODS

Lichen samples of four species were collected in spunbound olefin bags at various localities in different parts of both wilderness areas for laboratory analysis. Species collected and the substrates were Cladina rangiferina on soil, Cladina stygia on soil, Hypogymnia physodes on conifer branches, Evernia mesomorpha on conifer branches. These species were selected because they are

locally present in abundance and relatively easy to clean.

Five localities were selected for elemental analysis and are indicated on the map of collection localities and in the locality list (Appendix I). These localities are: Rocky Branch Trail at the pass over the ridge, Lows Bald Spot off Madison Gulf Trail, northeast of Mt. Crawford, Mt. Eisenhower, and Wamsutta Trail north of Mt. Washington. Ten to 20 grams of each species were collected at each locality.

Lichens were air dried and cleaned of all bark and detritis under a dissecting microscope but thalli were not washed. Three samples of each collection were submitted for analysis. Some replicates of each species were ground before being divided for analysis and are so marked in the tables. Analysis was done for sulfur and multi-element analysis by the Research Analytical Laboratory at the University of Minnesota. In the sulfur analysis a ground and pelleted 100-150 mg sample was prepared for total sulfur by dry combustion and measurement of evolved sulfur dioxide on a LECO Sulfur Determinator, model no. SC-132, by infra red absorption. Multi-element determination for Ca, Mg, Na, K, P, Fe, Mn, Al, Cu, Zn, Cd, Cr, Ni, Pb, and B were determined simultaneously by Inductively Coupled Plasma (ICP) Atomic Emission Spectrometry. For the ICP one gram of dried plant material was dry ashed in a 20 ml high form silica crucible at 485 degrees Celsius for 10-12 hrs. Crucibles were covered during the ashing as a precaution against contamination. The dry ash was boiled in 2N HCl to improve the recovery of Fe, Al and Cr and followed by transfer of the supernatant to 7 ml plastic disposable tubes for direct determination by ICP.

RESULTS AND DISCUSSION

Table 1 gives the results of the analyses for all three replicates arranged by species. Table 2 gives the means and standard deviations for each set of replicates. Some of the reported values are below the lower detection limits of the instruments. If one reading was below the detection limit (indicated by * in the tables) 0.7 of the detection limit was used for that reading in the calculations. If two or more readings were below the detection limits (indicated by # in the tables) no calculations were done for that element on that species at that locality.

Table 1. Analysis of White Mt. Lichens
Values in ppm of thallus dry weight

Species	P	K	Ca	Mg	Al	Fe	Na	Mn	Zn	Cu	B	Pb	Ni	Cr	Cd	S	Locality
<u>Cladina rangiferina</u>	498	1507	494	257	109	102	21.7	197.7	13.3	1.6	0.6	2.0	1.0	0.3	#	460	Rky Br. Ridge
<u>Cladina rangiferina</u>	440	1315	421	226	96	89	27.7	179.0	12.2	1.6	0.5	3.7	0.9	0.3	#	450	Rky Br. Ridge
<u>Cladina rangiferina</u>	460	1364	479	244	103	99	26.9	149.2	12.9	1.6	0.6	2.6	0.8	0.2	#	430	Rky Br. Ridge
<u>Cladina rangiferina</u>	368	1464	347	209	128	139	25.8	24.3	19.6	2.0	0.6	10.5	0.9	0.3	0.2	530	Lows Bald Spot @
<u>Cladina rangiferina</u>	353	1458	339	203	129	135	26.7	24.1	20.6	2.0	0.6	8.7	1.1	0.2	0.1	560	Lows Bald Spot @
<u>Cladina rangiferina</u>	344	1468	341	207	128	135	24.3	24.3	18.9	2.0	0.6	8.0	1.1	0.4	0.2	590	Lows Bald Spot @
<u>Cladina rangiferina</u>	259	1014	245	159	197	216	29.2	59.6	16.5	1.9	0.6	10.5	1.0	0.5	0.1	510	NE Mt. Crawford
<u>Cladina rangiferina</u>	270	1060	258	170	165	175	27.1	52.3	17.6	1.8	0.5	8.9	0.6	0.3	0.2	440	NE Mt. Crawford
<u>Cladina rangiferina</u>	267	1003	249	159	165	171	32.9	96.4	15.7	1.7	0.5	7.8	1.0	0.3	0.2	450	NE Mt. Crawford
<u>Cladina rangiferina</u>	658	1963	445	240	149	195	23.0	86.9	29.3	2.0	0.7	6.7	1.0	0.4	0.3	590	Mt. Eisenhower
<u>Cladina rangiferina</u>	752	2105	459	252	145	181	18.4	92.2	30.5	2.2	0.6	5.8	1.1	0.4	0.2	600	Mt. Eisenhower
<u>Cladina rangiferina</u>	708	1917	466	242	142	184	16.9	96.6	32.1	2.1	0.6	5.3	1.0	0.5	0.2	640	Mt. Eisenhower
<u>Cladina stygia</u>	419	1656	354	199	114	118	19.9	56.5	18.8	1.9	0.5	5.7	1.0	0.1	0.2	540	Wamsutta Tr.
<u>Cladina stygia</u>	423	1640	363	192	130	133	21.1	51.1	19.6	1.8	0.5	7.7	0.7	0.3	0.3	630	Wamsutta Tr.
<u>Cladina stygia</u>	468	1779	340	207	112	115	23.3	54.2	18.3	1.7	0.6	8.2	0.8	*0.1	0.2	570	Wamsutta Tr.
<u>Hypogymnia physodes</u>	868	3300	6842	498	289	322	23.5	309.5	107.6	4.3	0.7	49.2	2.4	0.9	0.9	890	Rky Br. Ridge
<u>Hypogymnia physodes</u>	688	2544	9866	447	373	444	22.8	270.8	107.5	4.8	1.0	61.2	2.2	0.9	1.3	960	Rky Br. Ridge
<u>Hypogymnia physodes</u>	708	2896	5318	459	377	430	30.1	241.7	98.3	4.8	1.0	66.9	2.3	1.1	1.1	1030	Rky Br. Ridge
<u>Hypogymnia physodes</u>	791	2950	5981	507	327	393	34.3	170.3	97.7	4.2	1.2	51.9	3.2	0.8	0.6	1000	Lows Bald Spot @
<u>Hypogymnia physodes</u>	740	2863	5325	507	315	392	26.1	145.9	92.0	4.3	1.0	50.6	2.8	0.9	0.7	1030	Lows Bald Spot @
<u>Hypogymnia physodes</u>	752	2856	4851	522	302	367	30.9	181.1	89.8	4.2	1.1	47.0	3.2	0.9	0.6	1110	Lows Bald Spot @
<u>Hypogymnia physodes</u>	563	2383	6894	417	328	464	33.1	361.6	99.9	4.8	1.4	61.5	2.7	1.1	0.6	1070	NE Mt. Crawford
<u>Hypogymnia physodes</u>	545	2260	9556	392	313	428	29.5	263.2	101.1	5.1	1.2	64.5	2.6	0.9	0.9	1110	NE Mt. Crawford
<u>Hypogymnia physodes</u>	595	2551	7651	423	312	417	29.3	320.2	108.6	5.1	1.4	64.2	3.5	1.0	0.7	1105	NE Mt. Crawford
<u>Hypogymnia physodes</u>	721	2621	5598	534	414	486	35.0	164.6	97.5	5.8	1.4	64.3	3.2	1.1	1.0	1000	Mt. Eisenhower
<u>Hypogymnia physodes</u>	986	3078	7474	621	398	477	35.0	252.0	95.0	6.2	1.4	61.0	3.6	1.0	1.0	1040	Mt. Eisenhower
<u>Hypogymnia physodes</u>	798	2921	8762	558	408	468	34.8	208.3	98.6	6.7	1.7	61.8	3.2	1.2	1.5	1080	Mt. Eisenhower
<u>Hypogymnia physodes</u>	643	2472	21905	603	226	267	18.1	409.1	97.7	3.5	1.1	42.6	2.3	0.8	1.7	840	Wamsutta Tr.
<u>Hypogymnia physodes</u>	658	2499	19032	664	263	289	17.8	380.1	105.5	3.4	1.1	38.1	2.4	1.0	1.4	980	Wamsutta Tr.
<u>Hypogymnia physodes</u>	728	2732	14871	587	284	319	19.7	306.0	93.3	3.9	1.4	47.0	1.5	0.9	1.2	930	Wamsutta Tr.
<u>Evernia mesomorpha</u>	424	1782	220	192	220	251	33.9	29.8	37.2	2.7	1.1	25.3	1.1	0.7	0.3	1130	NE Mt. Crawford
<u>Evernia mesomorpha</u>	390	1700	197	178	195	211	33.0	22.5	32.6	2.4	1.1	21.4	1.3	0.5	0.3	1000	NE Mt. Crawford
<u>Evernia mesomorpha</u>	418	1749	213	180	185	199	33.1	28.7	32.8	2.5	1.0	21.8	0.5	0.5	0.4	1250	NE Mt. Crawford

* = one value at or below detection limit; included as 0.7 of detection limit

= two or more values at or below detection limit; not included in calculations

@ = ground before dividing into replicates

Table 2. Summary of White Mt. Lichens
Values in ppm of thallus dry weight

Cladina rangiferina

	P	K	Ca	Mg	Al	Fe	Na	Mn	Zn	Cu	B	Pb	Ni	Cr	Cd	S	Locality
Mean	466	1395	465	243	103	97	25.4	175.3	12.8	1.6	0.6	2.7	0.9	0.3	#	447	Rky Br. Ridge
Std. dev.	29	99	38	15	7	7	3.3	24.4	0.6	<.1	<.1	0.9	0.1	0.1	#	15	Rky Br. Ridge
Mean	355	1463	342	206	128	136	25.6	24.2	19.7	2.0	0.6	9.1	1.0	0.3	0.2	560	Lows Bald Spot @
Std. dev.	12	5	4	3	<.1	3	1.2	0.1	0.8	<.1	<.1	1.3	0.1	0.1	0.1	30	Lows Bald Spot @
Mean	266	1026	251	163	176	187	29.7	69.4	16.6	1.8	0.5	9.1	0.9	0.4	0.2	467	NE Mt. Crawford
Std. dev.	6	30	7	6	19	25	2.9	23.7	0.9	0.1	<.1	1.3	0.2	0.1	<.1	38	NE Mt. Crawford
Mean	706	1995	457	245	145	187	19.4	91.9	30.6	2.1	0.6	5.9	1.0	0.5	0.2	610	Mt. Eisenhower
Std. dev.	47	98	11	6	4	7	3.2	4.9	1.4	0.1	<.1	0.7	0.1	0.1	<.1	26	Mt. Eisenhower

Cladina stygia

	P	K	Ca	Mg	Al	Fe	Na	Mn	Zn	Cu	B	Pb	Ni	Cr	Cd	S	Locality
Mean	437	1692	352	199	119	122	21.4	53.9	18.9	1.8	0.5	7.2	0.8	*0.2	0.2	580	Wamsutta Tr.
Std. dev.	27	76	12	7	10	10	1.7	2.7	0.7	0.1	0.1	1.3	0.2	0.1	<.1	46	Wamsutta Tr.

Evernia mesomorpha

	P	K	Ca	Mg	Al	Fe	Na	Mn	Zn	Cu	B	Pb	Ni	Cr	Cd	S	Locality
Mean	410	1744	210	183	200	220	33.3	27.0	34.2	2.6	1.1	22.8	1.0	0.6	0.4	1127	NE Mt. Crawford
Std. dev.	18	41	12	7	18	27	0.5	3.9	2.6	0.2	<.1	2.2	0.4	0.1	<.1	125	NE Mt. Crawford

Hypogymnia physodes

	P	K	Ca	Mg	Al	Fe	Na	Mn	Zn	Cu	B	Pb	Ni	Cr	Cd	S	Locality
Mean	755	2913	7342	468	346	399	25.5	274.0	104.4	4.6	0.9	59.1	2.3	0.9	1.1	960	Rky Br. Ridge
Std. dev.	99	378	2315	27	50	67	4.0	34.0	5.3	0.3	0.2	9.0	0.1	0.1	0.2	70	Rky Br. Ridge
Mean	761	2890	5386	512	315	384	30.4	165.8	93.2	4.3	1.1	49.8	3.1	0.9	0.7	1047	Lows Bald Spot @
Std. dev.	27	53	567	8	13	15	4.1	18.0	4.1	<.1	0.1	2.5	0.2	<.1	0.1	57	Lows Bald Spot @
Mean	568	2398	8034	410	318	436	30.6	315.0	103.2	5.0	1.3	63.4	2.9	1.0	0.7	1095	NE Mt. Crawford
Std. dev.	25	146	1372	16	9	25	2.2	49.4	4.7	0.2	0.1	1.7	0.5	0.1	0.1	22	NE Mt. Crawford
Mean	835	2873	7278	571	407	477	34.9	208.3	97.1	6.2	1.5	62.3	3.3	1.1	1.2	1040	Mt. Eisenhower
Std. dev.	136	232	1591	45	8	9	0.1	43.7	1.8	0.4	0.1	1.7	0.2	0.1	0.3	40	Mt. Eisenhower
Mean	676	2568	18603	618	258	291	18.5	365.0	98.8	3.6	1.2	42.6	2.1	0.9	1.4	917	Wamsutta Tr.
Std. dev.	45	143	3537	41	30	26	1.0	53.2	6.2	0.2	0.2	4.5	0.5	0.1	0.2	71	Wamsutta Tr.

* = one value at or below detection limit; included as 0.7 of detection limit

= two or more values at or below detection limit; not included in calculations

@ = ground before dividing into replicates

All of the elemental levels found in the lichens are within typical limits for similar lichens. All element levels are very similar to those found in the Boundary Waters Canoe Area in Minnesota (a known clean area). From these tables it can be seen that there is no consistent correlation between element levels and location in the area. Although one species may have somewhat higher levels of one replicate at one locality, the other species may have higher levels at another locality so there is no overall correlation between high element levels and any one locality.

The sulfur levels in lichens tested range from 430 to 1250 ppm for all samples and these values are near background levels as cited by Solberg (1967) Erdman & Gough (1977), Nieboer et al (1977) and Puckett & Finegan (1980) for other species of lichens. These levels are very close to those reported in the same species in the Boundary Waters Canoe Area (Wetmore, 1987). Levels may be as low as 200-300 in the arctic (Tomassini et al, 1976) while levels in polluted areas are 4300-5200 ppm (Seaward, 1973) or higher. Different species may accumulate different amounts of elements and this is evident when comparing sulfur levels of different species. Cladina rangiferina and Cladina stygia have lower levels than Hypogymnia physodes and Evernia mesomorpha. Even when taking these differences into account there is no clear trend in accumulated levels of sulfur.

These tables indicate that there are no air pollution problems in the wilderness areas that can be detected with these methods.

CONCLUSIONS

The lichen flora is quite diverse and includes most of the species previously reported from the White Mts. There is no impoverishment of the lichen flora in any part of the study areas. However, some of the species reported from the area that were not found may have been lost because of the regional degradation of the air quality. The elemental analyses shows no elevated levels of sulfur or other common pollutant elements so there is probably no local source causing damage to the lichen flora. There are several species present in the most sensitive category to sulfur dioxide and some of these are rare. This rarity seems to be due more to ecological and climatic conditions than pollution since these species are quite healthy when present. The maps of the distributions of the more sensitive species do not show any significant voids that are not due to

normal ecological conditions. There is no evidence of damaged or dead lichens in any area where healthy ones are not also present. The elemental analyses do not show abnormal accumulations of polluting elements at any locality.

LITERATURE CITED

- Ahti, T. 1962. Notes on the lichen *Cladonia pseudorangiformis* Asah. Arch. Soc. Zool.-Bot. Fenn. 'Vanamo' 17:38-41.
- Ahti, T. 1966. *Parmelia olivacea* and the allied non-isidiate and non-sorediate corticolous lichens in the Northern Hemisphere. Acta Bot. Fenn. 70:1-68.
- Ahti, T. 1980. Taxonomic revision of *Cladonia gracilis* and its allies. Ann. Bot. Fenn. 17:195-243.
- Berry, E. C. 1941. A monograph of the genus *Parmelia* in North America, north of Mexico. Ann. Missouri Bot. Garden 28: 31-146.
- Bliss, L. C. 1962. Caloric and lipid content in alpine tundra plants. Ecology 43:753-757.
- Bliss, L. C. 1963. Alpine plant communities of the Presidential Range, New Hampshire. Ecology 44:678-697.
- Bliss, L. C. 1966. Plant productivity in alpine microenvironments on Mt. Washington, New Hampshire. Ecological Monographs 36:125-155.
- Bliss, L. C. & E. B. Hadley, 1964. Photosynthesis and respiration of alpine lichens. Amer. Jour. Bot. 51:870-874.
- Bowler, P. A. & P. W. Rundel. 1974. The Ramalina intermedia complex in North America. Bryologist 77:617-623.
- Brodo, I. M. & D. L. Hawksworth. 1977. *Alectoria* and allied genera in North America. Opera Botanica 42: 1-164.
- Brodo, I. M. 1984. The North American species of the Lecanora subfucsa group. Nova Hedwigia, Beih. 79:63-185.
- Culberson, W. L. 1958. The chemical strains of the lichen Parmelia cetrariooides Del. in North America. Phyton 11:85-92.

- Culberson, W. L. 1961. The *Parmelia quercina* group in North America. Amer. Jour. Bot. 48:168-174.
- Culberson, W. L. 1962. Some pseudocyclophellate *Parmeliae*. Nova Hedwigia 4: 563-577.
- Culberson, W. L. 1963. A summary of the lichen genus *Haematomma* in North America. Bryologist 66:224-236.
- Culberson, W. L. 1965. *Cetraria chicitae*, a new and widely distributed lichen species. Bryologist 68: 95-99.
- Culberson, W. L. 1969. The chemistry and systematics of some species of the *Cladonia cariosa* group in North America. Bryologist 72:377-386.
- Culberson, W. L. & C. F. 1956. The systematics of the *Parmelia dubia* group in North America. Amer. Jour. Bot. 43:678-687.
- Culberson, W. L. & C. F. Culberson. 1968. The lichen genera *Cetrelia* and *Platismatia* (*Parmeliaceae*). Contrib. U. S. Nat. Herbarium 34 (7):449-558.
- Dibben, M. J. 1980. The chemosystematics of the lichen genus *Pertusaria* in North America north of Mexico. Milwaukee Public Museum, Biol. 5:1-162.
- Erdman, J. A. & L. P. Gough. 1977. Variation in the element content of *Parmelia chlorochroa* from the Powder River Basin of Wyoming and Montana. Bryologist 80:292-303.
- Esslinger, 1977b. Studies in the lichen family Physciaceae. I. A new North American species. Mycotaxon 5:299-306.
- Esslinger, T. L. 1977. A chemosystematic revision of the brown *Parmeliae*. Jour. Hattori Bot. Lab. 42:1-211.
- Esslinger, T. L. 1978. Studies in the lichen family Physciaceae. II. The genus *Phaeophyscia* in North America. Mycotaxon 7:283-320.
- Flint, W. F. 1874. The distribution of plants in New Hampshire. In Hitchcock, C. H. The geology of New Hampshire, Part I. Physical Geography. Pp. 381-415 (lichens pp. 413-414).
- Hadley, E. B. & L. C. Bliss, 1964. Energy relationships of alpine plants on Mt. Washington, New Hampshire. Ecological Monographs 34:331-357.

- Hale, M. E. 1955a. Xanthoparmelia in North America I. The *Parmelia conspersa*-*stenophylla* group. Bull. Torrey Bot. Club 82:9-21.
- Hale, M. E. 1955b. Studies on the chemistry and distribution of North American lichens. Bryologist 58:242-246.
- Hale, M. E. 1956. Studies on the chemistry and distribution of North American lichens (6-9). Bryologist 59:114-117.
- Hale, M. E. 1958. Studies on the chemistry and distribution of North American lichens (10-13). Bryologist 61:81-85.
- Hale, M. E. 1963. Populations of chemical strains in the lichen *Cetraria ciliaris*. Brittonia 15:126-133.
- Hale, M. E. 1964. The *Parmelia conspersa* group in North America and Europe. Bryologist 67:462-473.
- Hale, M. E. 1976. A monograph of the lichen genus *Parmelia* Hasle (Parmeliaceae). Smithsonian Contributions to Botany Number 33, 60pp.
- Henssen, A. 1963. The North American species of *Placynthium*. Can. Jour. Bot. 41:1687-1724.
- Hutchinson, T. C., M. Dixon & M. Scott. 1986. The effect of simulated acid rain on feather mosses and lichens of the boreal forest. Water, Air, and Soil Pollution 31: 409-416.
- Imshaug, H. A. 1957. The lichen genus *Pyxine* in North and Middle America. Trans. Amer. Microsc. Soc. 76:246-269.
- Imshaug, H. A. & I. M. Brodo. 1966. Biosystematic studies on *Lecanora pallida* and some related lichens in the Americas. Nova Hedwigia 12:1-59.
- Jordan, W. P. 1973. The genus *Lobaria* in North America north of Mexico. Bryologist 76:225-251.
- K"arnefelt, I. 1979. The brown fruticose species of *Cetraria*. Opera Botanica 46:1-150.
- Kurokawa, S. 1962. A monograph of the genus *Anaptychia*. Beih. Nova Hedw. 6:1-115.
- Lang, G. E., W. A. Reiners & R. K. Heier. 1976. Potential alteration of precipitation chemistry

by epiphytic lichens. *Oecologia* 25:229-241.

Lang, G. E., W. A. Reiners & L. W. Pike. 1980. Structure and biomass dynamics of epiphytic lichen communities of balsam fir forests in New Hampshire. *Ecology* 61:541-550.

Laundon, J. R. 1967. A study of the lichen flora of London. *Lichenologist* 3:277-327.

LeBlanc, F., D. N. Rao & G. Comeau. 1972. The epiphytic vegetation of *Populus balsamifera* and its significance as an air pollution indicator in Sudbury, Ontario. *Canadian Journal of Botany* 50:519-528.

Lechowicz, M. J. 1987. Resistance of the caribou lichen *Cladina stellaris* (Opiz.) Brodo to growth reduction by simulated acidic rain. *Water, Air, and Soil Pollution* 34:71-77.

Llano, G. A. 1950. A monograph of the lichen family Umbilicariaceae in the western hemisphere. Off. Naval Res., Washington, D. C. (Navexos P-831):1-281.

Marsh, J. E. & T. H. Nash III. 1979. Lichens in relation to the Four Corners power plant in New Mexico. *The Bryologist* 82: 20-28.

Medlin, J. 1985. Using lichens to monitor acid rain in Michigan. *Mich. Bot.* 24:71-75.

Nash, T. H., III. 1973. Sensitivity of lichens to sulfur dioxide. *The Bryologist* 76:333-339.

Nash, T. H. & M. R. Sommerfeld. 1981. Elemental concentrations in lichens in the area of the Four Corners Power Plant, New Mexico. *Envir. and Exp. Botany* 21:153-162.

Nieboer, E., H. M. Ahmed, K. J. Puckett & D. H. S. Richardson. 1972. Heavy metal content of lichens in relation to distance from a nickel smelter in Sudbury, Ontario. *Lichenologist* 5:292-304.

Nieboer, E., K. J. Puckett, D. H. S. Richardson, F. D. Tomassini & B. Grace. 1977. Ecological and physiochemical aspects of the accumulation of heavy metals and sulphur in lichens. International Conference on Heavy Metals in the Environment, Symposium Proceedings 2(1):331-352.

Nieboer, E., D. H. S. Richardson & F. D. Tomassini. 1978. Mineral uptake and release by lichens: An Overview. *Bryologist* 81:226-246.

Puckett, K. J. & E. J. Finegan. 1980. An analysis of the element content of lichens from the Northwest Territories, Canada. *Can. Jour. Bot.* 58:2073-2089.

Scott, M. G. & T. C. Hutchinson. 1987. Effects of a simulated acid rain episode on photosynthesis and recovery in the caribou-forage lichens, Cladina stellaris (Opiz.) Brodo and Cladina rangiferina (L.) Wigg. *New Phytol.* 107:567-575.

Seaward, M. R. D. 1973. Lichen ecology of the Scunthorpe heathlands I. Mineral accumulation. *Lichenol.* 5:423-433.

Showman, R. E. 1975. Lichens as indicators of air quality around a coal-fired power generating plant. *Bryologist* 78:1-6.

Sierk, H. A. 1964. The genus Leptogium in North America north of Mexico. *Bryologist* 67:245-317.

Sigal, L. & J. Johnston. 1986. The effects of simulated acid rain on one species each of Pseudoparmelia, Usnea, and Umbilicaria. *Water, Air, and Soil Pollution* 27:315-322.

Solberg, Y. J. 1967. Studies on the chemistry of lichens. IV. The chemical composition of some Norwegian lichen species. *Ann. Bot. Fenn.* 4:29-34.

Thomson, J. W. 1950. The species of *Peltigera* of North America north of Mexico. *Amer. Midl. Nat.* 44:1-68.

Thomson, J. W. 1963. The lichen genus *Physcia* in North America. *Nova Hedwigia, Beih.* 7, viii + 172.

Thomson, J. W. 1967. The lichen genus Baeomyces in North America north of Mexico. *Bryologist* 70:286-298.

Tomassini, F. D., K. J. Puckett, E. Nieboer, D. H. S. Richardson & B. Grace. 1976. Determination of copper, iron, nickel, and sulphur by X-ray fluorescence in lichens from the Mackenzie Valley, Northwest Territories, and the Sudbury District, Ontario. *Can. Jour. Bot.* 54:1591-1603.

Trass, H. 1973. Lichen sensitivity to air pollution and index of paleotolerance (I.P.). *Folia Cryptogamica Estonica, Tartu*, 3:19-22.

Tuckerman, E. 1839. An enumeration of some lichens of New England, with remarks. *Boston Jour. Nat. Hist.* 2:245-262.

Tuckerman, E. 1840. A further enumeration of some New England lichens. *Boston Jour. Nat.*

Hist. 3:281-306.

Tuckerman, E. 1841. Further notices of some New England lichens. Boston Jour. Nat. Hist. 3:438-464.

Tuckerman, E. 1845. A further enumeration of some alpine and other lichens of New England. Boston Jour. Nat. Hist. 5:93-104.

Tuckerman, E. 1847. A synopsis of the lichens of the northern United States and British America. Proc. Acad. Arts & Sci. 1:195-284.

Tuckerman, E. 1858. Supplement to an enumeration of North American lichens; Part first, containing brief diagnosis of new species. Amer. Jour. Sci. & Arts, ser. 2. 25:422-430.

Tuckerman, E. 1859. Supplement to an enumeration of North American lichens, continued. Amer. Jour. Sci. & Arts, ser. 2. 28:200-206.

Tuckerman, E. 1860. Observations on North American and some other lichens, 1. Proc. Amer. Acad. Arts & Sci. 4:383-407.

Tuckerman, E. 1862. Observations on North American and some other lichens, 2. Proc. Amer. Acad. Arts & Sci. 5:383-422.

Tuckerman, E. 1864. Observations on North American and some other lichens, 3. Proc. Amer. Acad. Arts & Sci. 6:263-287.

Tuckerman, E. 1877. Observations on North American and some other lichens, 4. Proc. Amer. Acad. Arts & Sci. 12:166-185.

Tuckerman, E. 1872. Genera Lichenum: an Arrangement of the North American Lichens. xv, 281 pp. Amherst, Mass.

Tuckerman, E. 1882. A synopsis of North American Lichens; Part I, Comprising the Parmeliacei, Cladoniei, and Coenogoniei. xx, 261 pp. Boston.

Tuckerman, E. 1888. A synopsis of North American Lichens: II., Comprising the Lecideacei, and (in part) the Graphidacei. 176 pp. New Bedford, Mass.

Wetmore, C. M. 1960. The lichen genus Nephroma in North and Middle America. Publ. Museum, Michigan State Univ., Biol. Ser. 1:369-452.

Wetmore, C. M. 1982. Lichen decomposition in a black spruce bog. *Lichenologist* 14:267-271.

Wetmore, C. M. 1983. Lichens of the Air Quality Class 1 National Parks. Final Report, submitted to National Park Service, Air Quality Division, Denver, Colo.

Wetmore, C. M. 1987. Lichens and air quality in Boundary Waters Canoe Area of the Superior National Forest. Report submitted to U. S. Forest Service, Duluth.

APPENDIX I

Collection Localities

Collection numbers are those of Clifford Wetmore. All collections are listed in ascending order by collection number and date of collection.

Coos County

61580- Great Gulf Wilderness Area. NE slope of Mt. Clay. In 61649 alpine area with rock talus, heath meadow and some areas of krumholz with balsam fir, spruce and birch, elev. 6050 ft. 7 July 1988.

61650- Great Gulf Wilderness Area. Madison Gulf Trail at 61719 stream crossing about 1 mile from road. Along stream with balsam fir, black spruce and birch, elev. 1550 ft. 8 July 1988.

61720- Great Gulf Wilderness Area. Lows Bald Spot off Madison 61777 Gulf Trail near Mt. Washington Auto Road. Rocky knob with small balsam fir and black spruce, elev. 2875 ft. 8 July 1988. CHEMICAL ANALYSIS.

61778- Great Gulf Wilderness Area. Around Great Gulf Trail and 61849 W. Peabody River. Along ridge and river with black spruce, some balsam fir and rocks, elev. 2300 ft. 9 July 1988.

61850- Great Gulf Wilderness Area. Near W. Peabody River and 61891 Great Gulf Trail at NE corner of wilderness area. Back from stream with sugar maple, yellow birch, balsam fir and some black spruce, elev. 1725 ft. 9 July 1988.

61892- Presidential Dry River Wilderness Area. Knob half mile 61946 NE of Mt. Crawford along Davis Path. Rocky hilltop with black spruce and some balsam fir, elev. 2950 ft. 10 July 1988. CHEMICAL ANALYSIS.

61947- Presidential Dry River Wilderness Area. One mile SW of 61979 Mt. Crawford along Davis Path. At edge of ridge with balsam fir, birch and black spruce, elev. 2600 ft. 10 July 1988.

61980- Presidential Dry River Wilderness Area. Rocky Branch 62015 Trail at junction with Rocky Branch River E of Mt Davis. Along stream in balsam fir, white birch and mountain ash, elev. 2900 ft. 12 July 1988.

62016- Presidential Dry River Wilderness Area. North of 62052 Rocky Branch Trail at pass over ridge, E. of Mt. Davis. On rocky ridge just below peak with balsam fir, elev. 3450 ft. 12 July 1988. CHEMICAL ANALYSIS.

62053- Presidential Dry River Wilderness Area. Stairs Col 62082 Trail along Lower Stairs Brook NE of Stairs Mt. In valley along stream with yellow birch, beech and sugar maple, elev. 2100 ft. 13 July 1988.

Carroll County

62083- Presidential Dry River Wilderness Area. Along Rocky 62136 Branch River and trail NE of Maple Mt. At bend in river with birch, maple beech, balsam fir and some hemlock, elev. 1800 ft. 13 July 1988.

Coos County

62137- Presidential Dry River Wilderness Area. Along Dry 62180 River 1.5 miles from highway 302 at bridge. In river bottom with red maple, yellow birch and few beech and spruce, elev. 1700 ft. 14 July 1988.

62181- Great Gulf Wilderness Area. Wamsutta Trail north of 62243 Mt. Washington auto road. Around rocky knob 0.75 miles N of road with stunted balsam fir and black spruce and rocks, elev. 4350 ft. 15 July

1988. CHEMICAL ANALYSIS.

62244- Great Gulf Wilderness Area. Wamsutta Trail just below 62285 Mt. Washington auto road. In alpine meadow with rocks, heath and stunted balsam fir, elev. 5000 ft. 15 July 1988.

62286- Great Gulf Wilderness Area. Below Halfway House site 62327 on Mt. Washington auto road. On steep slope with balsam fir and yellow birch, elev. 3600 ft. 16 July 1988.

62328- Presidential Dry River Wilderness Area. Mt. Monroe on 62378 SW side in patches of balsam fir krumholz and openings, elev. 5000 ft. 17 July 1988.

62379- Presidential Dry River Wilderness Area. On east side 62428 of Mt. Monroe in sedge and heath meadow in alpine area, elev. 5000 ft. 17 July 1988.

62429- Presidential Dry River Wilderness Area. South side of 62462 Mt. Eisenhower (Mt. Pleasant of Tuckerman). In dwarf balsam fir along trail on hillside, elev. 4400 ft. 18 July 1988. CHEMICAL ANALYSIS.

62463- Presidential Dry River Wilderness Area. Mt. Franklin. 62504 Near top on south side in alpine meadow with rocks, elev. 4900 ft. 18 July 1988.

62505 White Mts. Nat. Forest. On concrete at Lakes of the Clouds Hut near Mt. Monroe, elev. 5000 ft. 18 July 1988.

62506- Presidential Dry River Wilderness Area. Upper end of 62541 Oakes Gulf near Camel Trail. In alpine meadow of sedge and heath and rocks, elev. 5350 ft. 19 July 1988.

62542- Presidential Dry River Wilderness Area. Two miles up 62584 Dry River and above Clinton Trail junction. Along stream with yellow birch, red maple, balsam fir and some black spruce, elev. 2200 ft. 21 July 1988.

APPENDIX II

Species Sensitive to Sulfur Dioxide

Based on the list of lichens with known sulfur dioxide sensitivity compiled from the literature, the following species in the White Mt. wilderness areas are within the Sensitive category as listed by Wetmore, 1983. Sensitive species (S) are those present only under 50ug sulfur dioxide per cubic meter (average annual). The intermediate category includes species present between 50ug and 100ug. The SI group falls between the Sensitive and Intermediate categories. Open circles on the maps are localities where the species was not found and solid circles are where it was found. Note: Refer to text for interpretation of these maps and precautions concerning absence in parts of the park.

Fig. 2 Bryoria furcellata

Fig. 3 Lecidea vernalis

Fig. 4 Lobaria pulmonaria

Fig. 5 Ochrolechia androgyna

Fig. 6 Parmelia squarrosa

Fig. 7 Usnea filipendula

APPENDIX III

Lichens reported in the literature from the White Mts.

Names have been brought up to date with present nomenclature whenever possible. In some cases the old names were missapplied and without studying the original collections it is not possible to determine what the present name should be. In this case the old name is enclosed in brackets.

- [Acarospora cervina Tuckerman, 1847, Flint, 1874 Misident.]
Acarospora sinopica (Wahlenb. in Ach.) K"orb. Tuckerman, 1882
Alectoria fallacina Mot. Brodo & Hawksworth, 1977
[Alectoria ochroleuca (Hoffm.) Mass. Flint, 1874 Misident.]
Alectoria sarmentosa (Ach.) Ach. Tuckerman, 1845, Brodo & Hawksworth, 1977
Amygdalaria panaeola (Ach.) Hertel & Brodo Tuckerman, 1847, 1888, Flint, 1874
Anaptychia palmulata (Michx.) Vain. Tuckerman, 1845, Kurokawa, 1962, Hale, 1956
Anzia colpodes (Ach.) Stizenb. Hale, 1955b
Arthonia cinereopruinosa Schaer. Tuckerman, 1872, Flint, 1874
Arthonia diffusa Nyl. Tuckerman, 1872, Flint, 1874
[Arthonia luridoalba Nyl. Flint, 1874 Misident.]
Arthonia patellulata Nyl. Tuckerman, 1872, Flint, 1874
Arthonia spadicea Leight. Tuckerman, 1872, Flint, 1874
Aspicilia cinerea (L.) K"orb. Tuckerman, 1840
Aspicilia gibbosa (Ach.) K"orb. Tuckerman, 1882
Bacidia bagleittona (Mass. & De Not. in Mass.) Jatta Tuckerman, 1840, 1872, 1888
Bacidia chlorantha (Tuck.) Fink Tuckerman, 1872, 1888, Flint, 1874
Bacidia inundata (Fr.) K"orb. Tuckerman, 1888
[Bacidia sabuletorum (Schreb.) Lett. Flint, 1874 Misident.]
Bacidia sphaeroides (Dicks.) Zahlbr. Tuckerman, 1888
Baeomyces placophyllus Ach. Tuckerman, 1845, 1847, 1888, Flint, 1874, Thomson, 1967
Baeomyces roseus Pers. Tuckerman, 1847, Flint, 1874, Thomson, 1967
Baeomyces rufus (Huds.) Rebent. Tuckerman, 1839, 1847, 1888, Flint, 1874, Thomson, 1967
Botrydina viridis (Ach.) Redh. & Kuyper Tuckerman, 1847, 1872, Flint, 1874
Bryoria bicolor (Ehrh.) Brodo & Hawksw. Tuckerman, 1840, 1847, 1882, Brodo & Hawksworth, 1977
Bryoria chalybeiformis (L.) Brodo & Hawksw. Tuckerman, 1839, 1882
Bryoria furcellata (Fr.) Brodo & Hawksw. Brodo & Hawksworth, 1977
Bryoria fuscescens (Gyeln.) Brodo & Hawksw. Brodo & Hawksworth, 1977
Bryoria nadvornikiana (Gyeln.) Brodo & Hawksw. Brodo & Hawksworth, 1977, Lang et al., 1980
Bryoria tenuis (Dahl) Brodo & Hawksw. Brodo & Hawksworth, 1977
Bryoria trichodes (Michx.) Brodo & Hawksw. Brodo & Hawksworth, 1977
Buellia coracina (Hoffm.) K"orb. Tuckerman, 1888, Flint, 1874
Buellia schaeferi De Not. Tuckerman, 1872, 1888, Flint, 1874
Buellia wahlenbergii (Ach.) Sheard Tuckerman, 1872, 1888, Flint, 1874
Calicium abietinum Pers. Tuckerman, 1872, Flint, 1874
[Calicium adpersum Pers. Flint, 1874 Misident.]
Calicium salicinum Pers. Tuckerman, 1845, 1872, Flint, 1874
Caloplaca flavovirescens (Wulf.) Dalla Torre & Sarnth. Tuckerman, 1847, 1888, Flint, 1874
Caloplaca saxicola (Hoffm.) Nordin Tuckerman, 1845
Catillaria globulosa (Fl"orke) Th. Fr. Tuckerman, 1872, 1888, Flint, 1874

- Catillaria griffithii (Sm.) Malme Tuckerman, 1872, Flint, 1874
Cetraria arenaria K"arnef. K"arnefelt, 1979
Cetraria ciliaris Ach. Tuckerman, 1841, Hale, 1963, Lang et al., 1980
Cetraria commixta (Nyl.) Th. Fr. Tuckerman, 1840, 1847, 1882, Flint, 1874
Cetraria cucullata (Bellardi) Ach. Tuckerman, 1839, 1840, 1847, 1882, Flint, 1874, Bliss, 1962, 1963
Cetraria delisei (Bory ex Schaer.) Nyl. K"arnefelt, 1979, Bliss, 1963
Cetraria ericetorum Opiz Tuckerman, 1840
Cetraria islandica (L.) Ach. Tuckerman, 1839, 1840, 1882, Bliss, 1962, 1963, 1966, Hadley & Bliss, 1964, Bliss & Hadley, 1964,
Cetraria islandica subsp. crispiformis (R"as.) K"arnef. K"arnefelt, 1979
Cetraria laevigata Rass. K"arnefelt, 1979
Cetraria nivalis (L.) Ach. Tuckerman, 1839, 1845, 1847, 1882, Flint, 1874, Bliss, 1962, 1963, Bliss & Hadley, 1964
Cetraria oakesiana Tuck. Tuckerman, 1841, Hale, 1956, Lang et al., 1980
Cetraria pinastri (Scop.) Gray Tuckerman, 1839, 1840, Lang et al., 1980
Cetraria sepincola (Ehrh.) Ach. Tuckerman, 1847, 1882, Flint, 1874, Lang et al., 1980
Cetrelia cetrariooides (Del. ex Duby) W. & C. Culb. Berry, 1941, Culberson, 1958, Culberson, 1962
Cetrelia chicitae (W. Culb.) W. & C. Culb. Culberson, 1965, Culberson & Culberson, 1968
Cetrelia olivetorum (Nyl.) W. & C. Culb. Culberson, 1962, Culberson & Culberson, 1968
[Chaenotheca ferruginea (Turn. ex Sm.) Mig. Flint, 1874 Misident.]
Chaenotheca phaeocephala (Turn.) Th. Fr. Tuckerman, 1872
Cladina mitis (Sandst.) Hustich Bliss, 1962, 1963
Cladina rangiferina (L.) Nyl. Bliss, 1962, 1963, 1966, Bliss & Hadley, 1964, Hadley & Bliss, 1964
Cladina stellaris (Opiz) Brodo Bliss, 1962, 1963
Cladonia amaurocraea (Fl"orke) Schaer. Tuckerman, 1847, 1882, Flint, 1874, Bliss, 1963
Cladonia bellidiflora (Ach.) Schaer. Tuckerman, 1847, 1882, Flint, 1874
Cladonia boryi Tuck. Tuckerman, 1841, 1847, 1882, Flint, 1874
Cladonia caespiticia (Pers.) Fl"orke Tuckerman, 1845
Cladonia cenotea (Ach.) Schaer. Tuckerman, 1847, 1882, Flint, 1874
Cladonia cervicornis (Ach.) Flot. Tuckerman, 1882
Cladonia cornuta (L.) Hoffm. Tuckerman, 1882
Cladonia cristatella Tuck. Tuckerman, 1858, 1862, 1882
Cladonia cyanipes (Somm.) Nyl. Tuckerman, 1847, Flint, 1874
Cladonia decorticata (Fl"orke) Spreng. Tuckerman, 1847, 1882, Flint, 1874
Cladonia digitata (L.) Hoffm. Tuckerman, 1840
Cladonia fimbriata (L.) Fr. Tuckerman, 1845, 1882, Flint, 1874
Cladonia floerkeana (Fr.) Fl"orke Bliss, 1963
Cladonia furcata (Huds.) Schrad. Tuckerman, 1839, 1882
Cladonia gracilis (L.) Willd. Tuckerman, 1839, 1840, 1845, Flint, 1874, Bliss, 1962, 1963
Cladonia gracilis subsp. nigripes (Nyl.) Ahti Tuckerman, 1840
Cladonia macilenta Hoffm. Tuckerman, 1841, 1882
Cladonia maxima (Asah.) Ahti Ahti, 1980
Cladonia parasitica (Hoffm.) Hoffm. Tuckerman, 1845
Cladonia pseudorangiformis Asah. Ahti, 1962
[Cladonia pyxidata (L.) Hoffm. Flint, 1874 Misident.]
Cladonia squamosa (Scop.) Hoffm. Tuckerman, 1840, Lang et al., 1980
Cladonia uncialis (L.) Web. ex Wigg. Bliss, 1963, Hadley & Bliss, 1964
Cladonia verticillata (Hoffm.) Schaer. Tuckerman, 1882
Cornicularia aculeata (Schreb.) Ach. Tuckerman, 1840, 1847, 1882, Flint, 1874, Bliss, 1963
Dendriscocaulon intricatum (Nyl.) Hanssen Tuckerman, 1872, 1882, Flint, 1874
Dirinaria frostii (Tuck.) Hale & W. Culb. Thomson, 1963
Eopyrenula leucoplaca (Wallr.) R. Harris Tuckerman, 1872, Flint, 1874
Ephebe lanata (L.) Vain. Tuckerman, 1845, 1847
Evernia mesomorpha Nyl. Tuckerman, 1847, Lang et al., 1976, Lang et al., 1980

- [Evernia purpuracea Flint, 1874 Unknown name]
- Fuscidea cyathoides (Ach.) V. Wirth & Vezda Tuckerman, 1845, Flint, 1874
- Fuscidea mollis (Wahlenb.) V. Wirth & Vezda Tuckerman, 1872
- Haematomma lapponicum R"as. Tuckerman, 1839, 1840, 1847, 1882, Flint, 1874, Culberson, 1963
- Haematomma ochrophaeum (Tuck.) Mass. Flint, 1874
- Heterodermia hypoleuca (Muhl.) Trevis. Tuckerman, 1845
- Hydrothyria venosa J. Russell Tuckerman, 1872, Flint, 1874
- Hypogymnia krogiae Ohls. Tuckerman, 1840, Berry, 1941, Lang et al., 1980
- Hypogymnia physodes (L.) Nyl. Berry, 1941, Lang et al., 1980
- Hypogymnia tubulosa (Schaer.) Hav. Lang et al., 1980
- Hypogymnia vittata (Ach.) Gas. Tuckerman, 1882, Lang et al., 1980
- Icmadophila ericetorum (L.) Zahlbr. Flint, 1874
- Ionaspis odora (Ach.) Th. Fr. in B. Stein Tuckerman, 1882
- Imshaugia aleurites (Ach.) S. F. Meyer Tuckerman, 1845, 1872, Flint, 1874, Lang et al., 1980
- Julella lactea (Mass.) Zahlbr. Tuckerman, 1872, Flint, 1874
- Lasallia papulosa (Ach.) Llano Tuckerman, 1839, Flint, 1874, Llano, 1950
- Lasallia pensylvanica (Hoffm.) Llano Tuckerman, 1839, Llano, 1950
- Lasallia pustulata (L.) Merat Tuckerman, 1845, 1882
- Lecanactis chloroconia Tuck. Tuckerman, 1864
- Lecania cyrtella (Ach.) Th. Fr. Tuckerman, 1872, Flint, 1874
- Lecanora allophana Nyl. Brodo, 1984
- Lecanora argentata (Ach.) Malme Brodo, 1984
- Lecanora argopholis (Ach.) Ach. Tuckerman, 1882
- Lecanora atriseda (Fr.) Nyl. Tuckerman, 1882
- Lecanora badia (Hoffm.) Ach. Tuckerman, 1882, Flint, 1874
- Lecanora caesiorubella Ach. subsp. caesiorubella Imshaug & Brodo, 1966
- Lecanora caesiorubella subsp. prolifera (Fink) R. Harris Imshaug & Brodo, 1966
- Lecanora cinereofusca Magn. Brodo, 1984
- Lecanora circumborealis Brodo & Vitik. Brodo, 1984
- [Lecanora discreta Flint, 1874 Misident.]
- Lecanora glabrata (Ach.) Malme Brodo, 1984
- Lecanora granatina Somm. Tuckerman, 1872, 1882, Flint, 1874
- Lecanora hybocarpa (Tuck.) Brodo Brodo, 1984
- Lecanora intricata (Ach.) Ach. Tuckerman, 1882
- Lecanora polytropa (Hoffm.) Rabenh. Tuckerman, 1841, 1882
- Lecanora straminea Wahlenb. ex Ach. Tuckerman, 1845
- Lecidea aglaea Somm. Tuckerman, 1872, 1888, Flint, 1874
- Lecidea auriculata Th. Fr. Tuckerman, 1888
- Lecidea caesioatra Schaer. Tuckerman, 1847, 1872, 1888, Flint, 1874
- Lecidea carnulenta (Tuck.) Fink Tuckerman, 1877, 1888
- [Lecidea confluens (Web.) Ach. Flint, 1874 Misident.]
- [Lecidea contigua b hydropheila Tuckerman, 1888 Name unknown]
- Lecidea diapensiae Th. Fr. Tuckerman, 1872, 1888, Flint, 1874
- Lecidea elabens Fr. Tuckerman, 1847, Flint, 1874
- Lecidea fuscoatra (L.) Ach. Tuckerman, 1888
- Lecidea fuscocinerea Nyl. Tuckerman, 1888
- Lecidea lapicida (Ach.) Ach. Tuckerman, 1872, 1888, Flint, 1874
- Lecidea lugubris Somm. Tuckerman, 1872, 1888, Flint, 1874
- Lecidea peliaspis (Tuck.) Zahlbr. Tuckerman, 1888
- Lecidea pycnocarpa (K"orb.) Ohl. Tuckerman, 1888
- Lecidea tenebrosa Flot. Tuckerman, 1872, 1888, Flint, 1874
- Lecidea tornensis Nyl. Tuckerman, 1888
- Lecidoma demissum (Rustr.) G. Schneid. & Hert. Tuckerman, 1847, 1872, 1888, Flint, 1874

- Lepraria membranacea (Dicks.) Vain. Tuckerman, 1845, Lang et al., 1980
Lepraria neglecta (Nyl.) Lett. Tuckerman, 1888
Leprocaulon microscopicum (Vill.) Gams ex D. Hawksw. Tuckerman, 1847, Flint, 1874
Leptogium burgessii (L.) Mont. Tuckerman, 1872, 1882
Leptogium cyanescens (Rabenh.) K"orb. Sierk, 1964
Leptogium saturninum (Dicks.) Nyl. Sierk, 1964
Leptogium tenuissimum (Dicks.) K"orb. Sierk, 1964
[Leptogium tremelloides Tuckerman, 1840 Misident.]
[Letharia vulpina (L.) Hue Flint, 1874 Misident.]
Lobaria linita (Ach.) Rabenh. Tuckerman, 1882, Flint, 1874
Lobaria pulmonaria (L.) Hoffm. Tuckerman, 1839, 1840, 1847, Flint, 1874, Jordan, 1973
Lobaria quercizans Michx. Tuckerman, 1845, Jordan, 1973
Lopadium pezizoideum (Ach.) K"orb. Tuckerman, 1888, Flint, 1874, Lang et al., 1980
Megalospora tuberculosa (Fee) Sipman Tuckerman, 1872, 1888, Flint, 1874
Menegazzia terebrata (Hoffm.) Mass. Tuckerman, 1840, 1882, Hale, 1955b
Micarea melaena (Nyl.) Hedl. Tuckerman, 1888
Microcalicium arenarium (Hampe ex Mass.) Tibell Tuckerman, 1872, Flint, 1874
Microcalicium disseminatum (Ach.) Vain. Tuckerman, 1872, Flint, 1874
Mycoblastus sanguinarius (L.) Norm. Lang et al., 1980
Mycocalicium subtile (Pers.) Szat. Tuckerman, 1845
Nephroma arcticum (L.) Torss. Tuckerman, 1839, 1847, 1882, Flint, 1874, Wetmore, 1960
Nephroma bellum (Spreng.) Tuck. Wetmore, 1960
Nephroma helveticum Ach. Wetmore, 1960
Nephroma parile (Ach.) Ach. Tuckerman, 1840, 1847, Flint, 1874, Wetmore, 1960
Nephroma resupinatum (L.) Ach. Tuckerman, 1839, Wetmore, 1960
Opegrapha atra Pers. Tuckerman, 1845
Opegrapha microcycla Tuck. Tuckerman, 1858
Pannaria conoplea (Ach.) Bory Tuckerman, 1845, 1847
Pannaria pezizoides (Web.) Trev. Tuckerman, 1882
[Pannaria rubiginosa (Ach.) Bory Flint, 1874 Misident.]
Parmelia alpicola Th. Fr. Tuckerman, 1882
Parmelia aurulenta Tuck. Tuckerman, 1882, Berry, 1941, Hale, 1958
Parmelia caperata (L.) Ach. Berry, 1941
Parmelia centrifuga (L.) Ach. Tuckerman, 1845, 1882, Flint, 1874, Berry, 1941
Parmelia cetrata Ach. Berry, 1941
Parmelia coniocarpa Laurer Berry, 1941
Parmelia conspersa (Ach.) Ach. Berry, 1941, Hale, 1955a, 1964, Bliss, 1963
Parmelia exasperata De Not. Esslinger, 1977
Parmelia galbina Ach. Berry, 1941, Culberson, 1961, Hale, 1976
Parmelia glabratula (Lamy) Nyl. Esslinger, 1977
Parmelia halei Ahti Ahti, 1966, Esslinger, 1977
[Parmelia halseyana Tuck. Tuckerman, 1841 Name unknown]
Parmelia incurva (Pers.) Fr. Tuckerman, 1845, 1847, 1882, Flint, 1874, Berry, 1941
Parmelia livida Tayl. Culberson, 1961
Parmelia olivacea (L.) Ach. Berry, 1941
Parmelia omphalodes (L.) Ach. Tuckerman, 1882
Parmelia panniformis (Nyl.) Vain. Esslinger, 1977
Parmelia pulla Ach. Tuckerman, 1872, Flint, 1874
Parmelia revoluta Fl"orke Lang et al., 1980
Parmelia rupestris Ach. Berry, 1941, Culberson & Culberson, 1956
Parmelia saxatilis (L.) Ach. Tuckerman, 1882, Berry, 1941, Lang et al., 1976, Lang et al., 1980
Parmelia septentrionalis (Lynge) Ahti Ahti, 1966, Esslinger, 1977
Parmelia somloensis Gyeln. Hale, 1955a
Parmelia soreciata (Ach.) Th. Fr. Flint, 1874, Esslinger, 1977

- Parmelia sphaerospora Nyl. Tuckerman, 1847, Flint, 1874
Parmelia stygia (L.) Ach. Tuckerman, 1839, 1847, 1882, Flint, 1874, Berry, 1941, Esslinger, 1977
Parmelia subargentifera Nyl. Esslinger, 1977
Parmelia sulcata Tayl. Berry, 1941, Lang et al., 1980
Parmeliopsis ambigua (Wulf. in Jacq.) Nyl. Tuckerman, 1845, 1847, 1872, 1882, Flint, 1874, Lang et al., 1980
Parmeliopsis hyperopta (Ach.) Arn. Lang et al., 1980
Peltigera aphthosa (L.) Willd. Thomson, 1950
Peltigera canina (L.) Willd. Tuckerman, 1872, 1882, Flint, 1874, Thomson, 1950
Peltigera collina (Ach.) Schrad. Tuckerman, 1882
Peltigera didactyla (With.) Laundon Tuckerman, , 1872, 1882
Peltigera evansiana Gyeln. Thomson, 1950
Peltigera horizontalis (Huds.) Baumg. Thomson, 1950
Peltigera malacea (Ach.) Funck Tuckerman, 1845, 1882, Flint, 1874, Thomson, 1950
Peltigera membranacea (Ach.) Nyl. Thomson, 1950
Peltigera polydactyla (Neck.) Hoffm. Tuckerman, 1841, Thomson, 1950
Peltigera rufescens (Weis.) Humb. Thomson, 1950
Peltigera scabrosa Th. Fr. Tuckerman, 1882
Peltigera venosa (L.) Hoffm. Thomson, 1950
Pertusaria amara (Ach.) Nyl. Dibben, 1980
Pertusaria dactylina (Ach.) Nyl. Tuckerman, 1882, Dibben, 1980
Pertusaria globularis (Ach.) Tuck. Dibben, 1980
Pertusaria glomerata (Ach.) Schaer. Tuckerman, 1872, 1882, Flint, 1874
Pertusaria leucostoma (Bernh.) Mass. Dibben, 1980
Pertusaria macounii (Lamb) Dibb. Dibben, 1980
Pertusaria neoscotica Lamb Dibben, 1980
Pertusaria oculata (Dicks.) Th. Fr. Tuckerman, 1845, 1847, Flint, 1874
Pertusaria trachythallina Erichs. Dibben, 1980
Pertusaria velata (Turn.) Nyl. Dibben, 1980
Pertusaria waghornei Hult. Dibben, 1980
Phaeocalicium curtisii (Tuck.) Tibell Tuckerman, 1872, Flint, 1874
Phaeocalicium praecedens (Nyl.) A. Schmidt Tuckerman, 1888
Phaeophyscia chloantha (Ach.) Moberg Esslinger, 1978
Phaeophyscia ciliata (Hoffm.) Moberg Thomson, 1963, Esslinger, 1978, Flint, 1874
Phaeophyscia orbicularis (Neck.) Moberg Thomson, 1963
Phaeophyscia rubropulchra (Degel.) Moberg Thomson, 1963, Esslinger, 1977b
Phylliscum demangeonii (Moug. & Mont. in Mont.) Nyl. Tuckerman, 1872, 1882, Flint, 1874
Physcia aipolia (Ehrh. ex Humb.) F"urnr. Thomson, 1963
Physcia dubia (Hoffm.) Lett. Thomson, 1963
Physcia stellaris (L.) Nyl. Thomson, 1963
Physcia tenella (Scop.) DC. in Lam. & DC. Thomson, 1963
Physconia detersa (Nyl.) Poelt Thomson, 1963
Physconia muscigena (Ach.) Poelt Thomson, 1963
Pilophorus cereolus (Ach.) Th. Fr. in Hellb. Tuckerman, 1847, 1858, 1882, Flint, 1874
Placopsis gelida (L.) Linds. Tuckerman, 1882
Placynthiella uliginosa (Schrad.) Coppins & James Tuckerman, 1872, Flint, 1874
Placynthium flabellosum (Tuck.) Zahlbr. Tuckerman, 1882, Henssen, 1963
Platismatia glauca (L.) W. & C. Culb. Tuckerman, 1882, Culberson & Culberson, 1968, Lang et al., 1976, Lang et al., 1980
Platismatia tuckermanii (Oakes) W. & C. Culb. Culberson & Culberson, 1968, Lang et al., 1980
Polychidium muscicola (Sw.) Gray Tuckerman, 1872, 1882, Flint, 1874
Porina lectissima (Fr.) Zahlbr. Tuckerman, 1872, Flint, 1874
Porpidia flavocaerulescens (Hornem.) Hert. & Schwab Tuckerman, 1888
Porpidia macrocarpa (DC. in Lam. & DC.) Hert. & Schwab Tuckerman, 1888

- Porpidia speirea (Ach.) Kremp. Tuckerman, 1888
Pseudevernia cladonia (Tuck.) Hale & W. Culb. Tuckerman, 1882, Hale, 1955b, Lang et al., 1976,
Lang et al., 1980
Pseudevernia consocians (Vain.) Hale & W. Culb. Tuckerman, 1840, Berry, 1941, Hale, 1955b, Lang
et al., 1980
Pseudocypphellaria crocata (L.) Vain. Tuckerman, 1840
[Psora decipiens (Hedwig) Hoffm. Flint, 1874 Misident.]
Psoroma hypnorum (Vahl) Gray Tuckerman, 1847, 1882, Flint, 1874
Pycnothelia papillaria Duf. Tuckerman, 1840, 1847, 1882, Flint, 1874
Pyxine sorediata (Ach.) Mont. Tuckerman, 1845, Imshaug, 1957
[Ramalina calicaris (L.) Fr. Flint, 1874 Misident.]
Ramalina farinacea (L.) Ach. Tuckerman, 1841
Ramalina geniculata Hook. f. & Tayl. Tuckerman, 1882
Ramalina intermedia (Del. ex Nyl.) Nyl. Bowler & Rundel, 1974
Ramalina tenuis (Tuck.) G. K. Merr. Tuckerman, 1872
Rhizocarpon alpicola (Anzi) Rabenh. Tuckerman, 1872, 1888, Flint, 1874
Rhizocarpon badioatrum (Fl"ork ex Spreng.) Th. Fr. Tuckerman, 1888
Rhizocarpon geographicum (L.) DC. Tuckerman, 1840, Flint, 1874
Rhizocarpon obscuratum (Ach.) Mass. Tuckerman, 1839
Rhizocarpon oederi (Weber) K"orb. Tuckerman, 1841, 1888
Rinodina ascociscana Tuck. Tuckerman, 1858, 1872, 1882, Flint, 1874
Rinodina millaria Tuck. Tuckerman, 1872, 1888, Flint, 1874
Rinodina turfacea (Wahlenb.) K"orb. Tuckerman, 1872, 1882, Flint, 1874
Sarea difformis (Fr.) Fr. Tuckerman, 1888
Schismatomma pericleum (Ach.) Branth & Rostr. Tuckerman, 1888
Scoliciosporum chlorococcum (Graewe ex Sten.) Vezda Lang et al., 1980
Sphaerophorus fragilis (L.) Pers. Tuckerman, 1839, 1840, 1847, 1872, Flint, 1874
Sphaerophorus globosus (Huds.) Vain. Tuckerman, 1839, 1847, 1872, Flint, 1874
Sporastatia testudinea (Ach.) Mass. Tuckerman, 1847, 1872, 1888, Flint, 1874
[Staurothele fissa (Tayl.) Zw. Flint, 1874 Misident.]
Staurothele fuscocuprea (Nyl.) Zsch. Tuckerman, 1872
Stereocaulon condensatum Hoffm. Tuckerman, 1845, 1847, Flint, 1874
Stereocaulon dactylophyllum Fl"ork Tuckerman, 1845, 1847, Flint, 1874
Stereocaulon glaucescens Tuck. Tuckerman, 1840
Stereocaulon nanodes Tuck. Tuckerman, 1859, 1872, 1882, Flint, 1874
Stereocaulon paschale (L.) Hoffm. Tuckerman, 1840, 1845, 1847, Flint, 1874, Bliss, 1962, 1963
Stereocaulon tomentosum Fr. Tuckerman, 1847, Flint, 1874
Stereocaulon vesuvianum Pers. Tuckerman, 1845, Flint, 1874
Strigula stigmatella (Ach.) R. Harris Tuckerman, 1872, Flint, 1874
Thamnolia subuliformis (Ehrh.) W. Culb. Tuckerman, 1839, 1840
Thamnolia vermicularis (Sw.) Ach. ex Schaer. Tuckerman, 1882, Bliss, 1963
[Toninia caeruleonigrans (Lightf.) Th. Fr. Flint, 1874 Misident.]
Trapeliopsis flexuosa (Fr.) Coppins & James Tuckerman, 1888, Flint, 1874
Trapeliopsis gelatinosa Fl"ork Coppins & James Tuckerman, 1888
Trapeliopsis viridescens (Schrad.) Coppins & James Tuckerman, 1872, 1888, Flint, 1874
Trypethelium virens Tuck. ex Michen. in Darl. Tuckerman, 1872, Flint, 1874
Umbilicaria arctica (Ach.) Nyl. Tuckerman, 1839
Umbilicaria cylindrica (L.) Del. ex Duby Llano, 1950
Umbilicaria deusta (L.) Baumg. Tuckerman, 1839, 1840, 1882, Llano, 1950
Umbilicaria hirsuta (Sw. ex Westr.) Hoffm. Tuckerman, 1840, Flint, 1874
Umbilicaria hyperborea (Ach.) Hoffm. Tuckerman, 1840, 1847, 1882, Flint, 1874, Llano, 1950
Umbilicaria mammulata (Ach.) Tuck. Llano, 1950
Umbilicaria muehlenbergii (Ach.) Tuck. Tuckerman, 1840, 1847, 1882, Flint, 1874, Llano, 1950
Umbilicaria polyphylla (L.) Baumg. Tuckerman, 1840, 1847, 1882, Flint, 1874

Umbilicaria proboscidea (L.) Schrad. Tuckerman, 1839, 1840, 1847, 1882, Flint, 1874, Llano, 1950
Umbilicaria torrefacta (Lightf.) Schrad. Tuckerman, 1839, 1840, 1847, 1882, Flint, 1874, Llano, 1950,
Umbilicaria yellea (L.) Ach. Tuckerman, 1839, 1841, 1882, Llano, 1950
Usnea cavernosa Tuck. Tuckerman, 1882
Usnea longissima Ach. Tuckerman, 1845
Usnea trichodea Ach. Tuckerman, 1882
Verrucaria margacea (Wahlenb. in Ach.) Wahlenb. Tuckerman, 1872, Flint, 1874
Xylographa abietina (Pers.) Zahlbr. Tuckerman, 1872, 1888, Flint, 1874